Question Number: 57 Question Id: 640653576844 Question Type: MCQ Is Question

Mandatory: No Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction

Time: 0

**Correct Marks: 1** 

Question Label: Multiple Choice Question

Which among the following expressions would complete blank (v)?

**Options:** 

6406531926616. ✓ Speak up

6406531926617. **Speak out** 

6406531926618. \* Call back

## Sem1 Maths1

Yes

**Section Id:** 64065339023

Section Number: 3

Section type: Online

Mandatory or Optional: Mandatory

Number of Questions: 12

Number of Questions to be attempted: 12

Section Marks: 50

**Display Number Panel:** Yes

Group All Questions: No

**Enable Mark as Answered Mark for Review and** 

Clear Response :

Maximum Instruction Time:

Sub-Section Number: 1

**Sub-Section Id:** 64065382283

**Question Shuffling Allowed:** No

Is Section Default?: null

Question Number: 58 Question Id: 640653576845 Question Type: MCQ Is Question

Mandatory: No Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction

Time: 0

**Correct Marks: 0** 

Question Label: Multiple Choice Question

THIS IS QUESTION PAPER FOR THE SUBJECT "FOUNDATION LEVEL: SEMESTER 1:

MATHEMATICS FOR DATA SCIENCE I (COMPUTER BASED EXAM)"

ARE YOU SURE YOU HAVE TO WRITE EXAM FOR THIS SUBJECT?

CROSS CHECK YOUR HALL TICKET TO CONFIRM THE SUBJECTS TO BE WRITTEN.

(IF IT IS NOT THE CORRECT SUBJECT, PLS CHECK THE SECTION AT THE <u>TOP</u> FOR THE SUBJECTS REGISTERED BY YOU)

**Options:** 

6406531926619. VYES

6406531926620. \* NO

Question Number: 59 Question Id: 640653576846 Question Type: MCQ Is Question

Mandatory: No Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction

Time: 0

**Correct Marks: 0** 

Question Label: Multiple Choice Question

## **Instructions:**

- There are some questions which have functions with discrete valued domains (such as day, month, year etc).
- For NAT type question, enter only one right answer even if you get multiple answers for that particular question.
- Notations:
  - $\mathbb{R}$ = Set of real numbers
  - Q= Set of rational numbers
  - Z= Set of integers
  - N= Set of natural numbers
- The set of natural numbers includes 0.

### **Options:**

6406531926621. ✓ Useful Data has been mentioned above.

6406531926622. \* This data attachment is just for a reference & not for an evaluation.

Sub-Section Number: 2

**Sub-Section Id:** 64065382284

**Question Shuffling Allowed:** No

Is Section Default?: null

Question Id: 640653576847 Question Type: COMPREHENSION Sub Question Shuffling

Allowed : No Group Comprehension Questions : No Question Pattern Type : NonMatrix

Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

**Question Numbers : (60 to 62)** 

Question Label: Comprehension

Let  $S = \{Jan, Feb, March, April, May\}$  be a set of months of a particular year. Consider the following relations on the set S.

- $R_1 = \{ \text{ (Feb, Feb), (March, April), (April, March), (March, March) } \}$
- $R_2 = \{ (Jan, Jan), (Feb, Feb), (March, March), (April, April), (May, May) \}$

From the below list of given terms find out the best possible options for each of the given subquestions:

- 1) Reflexive relation
- 2) Symmetric relation
- 3) Transitive relation
- 4) Not a equivalence relation
- 5) Not a function
- 6) Injective (One-one) function
- 7) Surjective (Onto) Function

### **Sub questions**

Question Number: 60 Question Id: 640653576848 Question Type: SA Calculator: None

Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

**Correct Marks: 3** 

Question Label: Short Answer Question

 $R_1$  is \_\_\_\_\_\_\_. (Enter all correct options. Enter only the serial numbers of those options in increasing order without adding any comma or space in between them i.e., if your answer is 6 and  $R_1$  then you should enter  $R_2$ .

7, then you should enter 67]

**Response Type:** Numeric

**Evaluation Required For SA:** Yes

**Show Word Count:** Yes

**Answers Type:** Equal

**Text Areas :** PlainText

**Possible Answers:** 

245

Question Number: 61 Question Id: 640653576849 Question Type: SA Calculator: None

Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

Question Label : Short Answer Question	
$R_2$ is (Enter all correct options. Enter only	y the serial numbers of those options in
increasing order without adding any comma or space	e in between them i.e., if your answer is 3 and
4, then you should enter 34)	
Response Type: Numeric	
<b>Evaluation Required For SA :</b> Yes	
Show Word Count: Yes	
Answers Type: Equal	
Text Areas: PlainText	
Possible Answers :	
12367	
Question Number : 62 Question Id : 640653576850	Question Type : SA Calculator : None
Response Time: N.A Think Time: N.A Minimum In	struction Time : 0
Correct Marks : 2	
Question Label : Short Answer Question	
Find the cardinality of the set $(S \times S) \setminus R_2$ .	
Response Type: Numeric	
Evaluation Required For SA : Yes	
Show Word Count: Yes	
Answers Type: Equal	
Text Areas: PlainText	
Possible Answers :	
20	
Sub-Section Number :	3
Sub-Section Id :	64065382285
Question Shuffling Allowed :	Yes
Is Section Default? :	null

**Correct Marks: 3** 

Question Number: 63 Question Id: 640653576857 Question Type: MCQ Is Question

Mandatory: No Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction

Time: 0

**Correct Marks: 5** 

Question Label: Multiple Choice Question

Consider a polynomial  $p(x) = 0.3x^3(x^2 - 1)(x - 2)^2(x - 3)$ 

Which of the figure represents the polynomial p(x)?

## **Options:**





Figure 1

6406531926634. \*\*



6406531926635. \*\*

Figure 2





Figure 3



6406531926637. \*\*

Figure 4

Sub-Section Number: 4

**Sub-Section Id:** 64065382286

**Question Shuffling Allowed :** Yes

Is Section Default?: null

Question Number: 64 Question Id: 640653576856 Question Type: MSQ Is Question

Mandatory: No Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction

Time: 0

Correct Marks: 3 Max. Selectable Options: 0

Question Label: Multiple Select Question

Which of the following options is/are true?

## **Options:**

6406531926630.  $\blacksquare$  If m and n are x- intercept and y- intercept of the line 2x+3y=6, respectively, then m+n=6

6406531926631. \*\* The y- intercept of a line is the perpendicular distance to the line from the origin.

6406531926632.  $\checkmark$  The lines 2x + 3y = 6 and 3x - 2y = 6 are perpendicular to each other.

6406531926633.  $\checkmark$  The distance between two parallel lines 2x+3y=6 and 4x+6y=12 is 0.

**Sub-Section Number:** 5

**Sub-Section Id:** 64065382287

**Question Shuffling Allowed :** Yes

Is Section Default?: null

Question Number : 65 Question Id : 640653576863 Question Type : MSQ Is Question

Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time: 0

Correct Marks: 5 Max. Selectable Options: 0

Question Label: Multiple Select Question

Consider two polynomials  $p(x) = x^4 + 3x^3 - 9x + 8$  and  $q(x) = (x^2 + x)(x + 3)$ . Let r(x) be the remainder obtained when p(x) is divided by q(x). Let l(x) be the line that passes through the y- intercept and the minimum point in the graph of r(x), for reference follow the Figure: M1Q1-1.




Figure: M1Q1-1

Which of the following options is/are true?

# Options:

6406531926645. \* 
$$r(x) = x^2 - 6x + 9$$

6406531926646. 
$$\checkmark l(x) \equiv y = -3x + 8$$

6406531926647. **\*** 
$$l(x) \equiv y = -2x + 8$$

6406531926648.  $\checkmark$  The number of turning points in q(x) is 2

**Sub-Section Id:** 64065382288

**Question Shuffling Allowed :** Yes

Is Section Default?: null

Question Number: 66 Question Id: 640653576851 Question Type: SA Calculator: None

Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

**Correct Marks: 5** 

Question Label: Short Answer Question

A company opened recruitment for the post of data analyst. 500 candidates have applied for the post. 285 candidates are proficient in Python programming, 195 candidates are proficient in *C* programming, 115 candidates are proficient in Java programming, 45 candidates are proficient in Python and Java, 70 candidates are proficient in *C* and Python, 50 candidates are proficient in *C* and Java and 50 candidates don't know any of the programming languages. Find the number of candidates who are proficient in exactly one of the three programming languages.

Response Type: Numeric

**Evaluation Required For SA:** Yes

**Show Word Count:** Yes

**Answers Type:** Equal

**Text Areas:** PlainText

**Possible Answers:** 

325

Question Number: 67 Question Id: 640653576855 Question Type: SA Calculator: None

Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

**Correct Marks: 5** 

Question Label: Short Answer Question

Consider the points A(0,3), B(x,y), C(4,3), D(1,0) and E(3,1) in the coordinate system. Suppose the point B divides internally the line segment AC in the ratio k:1. If the area of triangle DEB is 2, then find the positive value of k.

**Response Type:** Numeric

**Evaluation Required For SA:** Yes

-12

Sub-Section Number: 7

**Sub-Section Id:** 64065382289

**Question Shuffling Allowed :** Yes

Is Section Default?: null

Question Number: 69 Question Id: 640653576861 Question Type: SA Calculator: None

Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

#### **Correct Marks: 3**

**Question Label: Short Answer Question** 

Consider a quadratic function  $q(x) = ax^2 + bx + c$ , where  $a, b, c \in \mathbb{R}$  and  $a \neq 0$  with the following information:

- The Maximum value attained by q(x) is at x = -1.
- Discriminant value of q(x) is 8.
- Slope of the function at x = 1 is 8.

Find the value q(2).

Response Type: Numeric

**Evaluation Required For SA:** Yes

**Show Word Count:** Yes

**Answers Type:** Equal

**Text Areas:** PlainText

**Possible Answers:** 

17

Sub-Section Number: 8

**Sub-Section Id:** 64065382290

**Question Shuffling Allowed:** No

Is Section Default?: null

Question Id: 640653576852 Question Type: COMPREHENSION Sub Question Shuffling

Allowed: No Group Comprehension Questions: No Question Pattern Type: NonMatrix

Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

**Question Numbers: (70 to 71)** 

Question Label: Comprehension

A company's profit function, given by  $P(x) = -2x^2 + 8x - 6$ , represents the relationship between the quantity (x) of the raw material used in the manufacturing process and the resulting profit. Use this information to answer the given subquestions.

### **Sub questions**

Question Number : 70 Question Id : 640653576853 Question Type : SA Calculator : None		
Response Time : N.A Think Time : N.A Minimum Instruction Time : 0		
Correct Marks : 2		
Question Label : Short Answer Question		
Find the quantity of the raw material such that the company has the maximum profit.		
Response Type: Numeric		
<b>Evaluation Required For SA :</b> Yes		
Show Word Count: Yes		
Answers Type: Equal		
Text Areas: PlainText		
Possible Answers :		
2		
Question Number : 71 Question Id : 640653576854 Question Type : SA Calculator : None		
Response Time : N.A Think Time : N.A Minimum Ins	truction Time : 0	
Correct Marks : 2		
Question Label : Short Answer Question		
Find the quantity of the raw material ( $x > 1$ ) such that the company has no profit.		
Response Type: Numeric		
Evaluation Required For SA: Yes		
Show Word Count: Yes		
Answers Type: Equal		
Text Areas: PlainText		
Possible Answers :		
3		
Sub-Section Number :	9	
Sub-Section Id :	64065382291	
Question Shuffling Allowed :	No	
Is Section Default? :	null	

Question Id: 640653576858 Question Type: COMPREHENSION Sub Question Shuffling

Allowed: No Group Comprehension Questions: No Question Pattern Type: NonMatrix

Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

**Question Numbers: (72 to 73)** 

Question Label: Comprehension

Consider a polynomial  $p(x) = (x^2 - 1)(x^3 - 4x^2 + 4x)$  $(x^3 - 10x^2 + 33x - 36)$  such that

- $t(x) = (x^3 6x^2 + 9x)$  divides p(x).
- $s(x) = (x^2 5x + 4)$  divides p(x).
- q(x) is the quotient when p(x) is divided by the polynomial z(x) = (x+1)t(x)s(x)

Use this information to answer the given subquestions:

### **Sub questions**

Question Number: 72 Question Id: 640653576859 Question Type: SA Calculator: None

Response Time: N.A Think Time: N.A Minimum Instruction Time: 0

**Correct Marks: 3** 

Question Label: Short Answer Question

If m is the number of distinct roots and n is the number of turning points of the polynomial p(x),

then find the value of m + n

Response Type: Numeric

**Evaluation Required For SA:** Yes

**Show Word Count:** Yes

**Answers Type:** Equal

**Text Areas:** PlainText

**Possible Answers:** 

13

Question Number: 73 Question Id: 640653576860 Question Type: MSQ Is Question

Mandatory: No Calculator: None Response Time: N.A Think Time: N.A Minimum Instruction

Time: 0

Correct Marks: 4 Max. Selectable Options: 0

Question Label: Multiple Select Question

Which of the following options is/are true?

**Options:** 

6406531926639.  $\checkmark$  The Minimum value of the quotient q(x) is 0.

6406531926640. \* p(x) is an odd degree polynomial.

6406531926641.  $\checkmark$  End behavior  $p(x) \to +\infty$  as  $x \to +\infty$ 

6406531926642.  $\checkmark$  End behavior  $p(x) \to +\infty$  as  $x \to -\infty$ 

## Sem1 Statistics1

**Section Id:** 64065339024

Section Number: 4

Section type: Online

Mandatory or Optional: Mandatory

Number of Questions: 10

Number of Questions to be attempted: 10

Section Marks: 40

**Display Number Panel:** Yes

**Group All Questions:** No

**Enable Mark as Answered Mark for Review and** 

Yes Clear Response:

**Maximum Instruction Time:** 0

Sub-Section Number: 1

**Sub-Section Id:** 64065382292