

**Show Word Count :** Yes

**Answers Type :** Equal

**Text Areas :** PlainText

**Possible Answers :**

90

**Question Number :** 164 **Question Id :** 640653588679 **Question Type :** SA **Calculator :** None

**Response Time :** N.A **Think Time :** N.A **Minimum Instruction Time :** 0

**Correct Marks :** 2

**Question Label :** Short Answer Question

Calculate the Herfindahl index. Round off to the nearest integer value between 0 and 10000

**Response Type :** Numeric

**Evaluation Required For SA :** Yes

**Show Word Count :** Yes

**Answers Type :** Equal

**Text Areas :** PlainText

**Possible Answers :**

2500

## MLT

<b>Section Id :</b>	64065339803
<b>Section Number :</b>	10
<b>Section type :</b>	Online
<b>Mandatory or Optional :</b>	Mandatory
<b>Number of Questions :</b>	13
<b>Number of Questions to be attempted :</b>	13
<b>Section Marks :</b>	50
<b>Display Number Panel :</b>	Yes
<b>Group All Questions :</b>	No

**Enable Mark as Answered Mark for Review and Clear Response :** Yes  
**Maximum Instruction Time :** 0  
**Sub-Section Number :** 1  
**Sub-Section Id :** 64065384980  
**Question Shuffling Allowed :** No  
**Is Section Default? :** null

**Question Number : 165 Question Id : 640653588681 Question Type : MCQ Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 0**

Question Label : Multiple Choice Question

**THIS IS QUESTION PAPER FOR THE SUBJECT "DIPLOMA LEVEL : MACHINE LEARNING TECHNIQUES (COMPUTER BASED EXAM)"**

**ARE YOU SURE YOU HAVE TO WRITE EXAM FOR THIS SUBJECT?**

**CROSS CHECK YOUR HALL TICKET TO CONFIRM THE SUBJECTS TO BE WRITTEN.**

**(IF IT IS NOT THE CORRECT SUBJECT, PLS CHECK THE SECTION AT THE [TOP](#) FOR THE SUBJECTS REGISTERED BY YOU)**

**Options :**

6406531963438. ✓ YES

6406531963439. ✗ NO

**Sub-Section Number :** 2  
**Sub-Section Id :** 64065384981  
**Question Shuffling Allowed :** Yes  
**Is Section Default? :** null

**Question Number : 166 Question Id : 640653588682 Question Type : MCQ Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction**

Time : 0

Correct Marks : 4

Question Label : Multiple Choice Question

Consider a training dataset of  $n$  points for a regression problem. Assume that the model is linear. Let  $\mathbf{w}_1$  and  $\mathbf{w}_2$  be the optimal weight vectors obtained from solving the following optimization problems.

$$\mathbf{w}_1 = \arg \min_{\mathbf{w}} \sum_{i=1}^n (\mathbf{w}^T \mathbf{x}_i - y_i)^2$$

$$\mathbf{w}_2 = \arg \min_{\mathbf{w}} \sum_{i=1}^n (\mathbf{w}^T \mathbf{x}_i - y_i)^3$$

Choose the most appropriate answer.

Options :

6406531963440. ✓  $\mathbf{w}_1$  will generalize better than  $\mathbf{w}_2$  on the test dataset.

6406531963441. ✗  $\mathbf{w}_2$  will generalize better than  $\mathbf{w}_1$  on the test dataset.

6406531963442. ✗ Both models will show identical performance on the test dataset.

Question Number : 167 Question Id : 640653588683 Question Type : MCQ Is Question

Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction

Time : 0

Correct Marks : 4

Question Label : Multiple Choice Question

The training dataset for a binary classification problem is as follows:

$$\{ (\mathbf{u}, 1), (-\mathbf{u}, 0), (2\mathbf{u}, 1), (-2\mathbf{u}, 0) \}$$

where,  $\mathbf{u} \in \mathbb{R}^d$  is a non zero constant and each element in the set given above is a data-point of the form  $(\mathbf{x}_i, y_i)$ . The labels lie in  $\{0, 1\}$ . Consider a linear classifier with weight vector  $\mathbf{w}$ . What condition should the weight vector satisfy for the zero-one loss to be zero on this dataset?

Options :

6406531963443. ✖  $\mathbf{w}^T \mathbf{u} < 0$

6406531963444. ✔  $\mathbf{w}^T \mathbf{u} > 0$

6406531963445. ✖  $\mathbf{w}^T \mathbf{u} = 0$

6406531963446. ✖ We can never find a  $\mathbf{w}$  for which the zero-one loss becomes zero on this dataset.

Sub-Section Number :	3
Sub-Section Id :	64065384982
Question Shuffling Allowed :	Yes
Is Section Default? :	null

Question Number : 168 Question Id : 640653588684 Question Type : MSQ Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0

Correct Marks : 4 Max. Selectable Options : 0

Question Label : Multiple Select Question

Consider a linear regression model that was trained on dataset  $X$  of shape  $(d, n)$ . Which of the following techniques could potentially decrease the loss on the training data (assuming the loss is the squared error)?

Options :

6406531963447. ✔ Adding a dummy feature in the dataset and learning the intercept  $w_0$  as well.

6406531963448. ✖ Penalizing the model weights with L2 regularization.

6406531963449. ✖ Penalizing the model weights with L1 regularization.

6406531963450.

✓ Training the kernel regression model of degree 2.

**Question Number : 169 Question Id : 640653588686 Question Type : MSQ Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 4 Max. Selectable Options : 0**

Question Label : Multiple Select Question

Which of the following statements are true about the decision tree algorithm?

**Options :**

6406531963455. ✖ Decision trees are prone to overfit if the maximum depth is set too low.

6406531963456. ✓ Decision trees are prone to underfit if the maximum depth is set too low.

6406531963457. ✓ Decision trees are sensitive to small perturbations in the dataset and can result in different tree structures.

6406531963458. ✓ Decision trees can handle both numerical and categorical features.

<b>Sub-Section Number :</b>	4
<b>Sub-Section Id :</b>	64065384983
<b>Question Shuffling Allowed :</b>	Yes
<b>Is Section Default? :</b>	null

**Question Number : 170 Question Id : 640653588685 Question Type : MSQ Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 4.5 Max. Selectable Options : 0**

Question Label : Multiple Select Question

Which of the following statements is/are true regarding solution of Ridge regression problem?

**Options :**

6406531963451. ✓ If there are multiple  $w$  solutions for minimizing mean square error, then  $w_R$  will be the one with least norm.

6406531963452. ✖ If there are multiple  $w$  solutions for minimizing mean square error, then  $w_R$  will be the one with highest norm.

6406531963453. ✔ Prior for  $w$  is  $N(0, \gamma^2 I)$  and  $y_i|x_i \sim N(w^T x_i, \sigma^2)$

6406531963454. ✖ Prior for  $w$  is  $N(1, \gamma^2 I)$  and  $y_i|x_i \sim N(0, \sigma^2)$

**Sub-Section Number :** 5  
**Sub-Section Id :** 64065384984  
**Question Shuffling Allowed :** Yes  
**Is Section Default? :** null

**Question Number : 171 Question Id : 640653588687 Question Type : SA Calculator : None**  
**Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**  
**Correct Marks : 4.5**

Question Label : Short Answer Question

Consider kernel regression with the kernel function  $(\mathbf{x}_1^T \mathbf{x}_2 + 2)^2$  applied on the following dataset.

$$\mathbf{X} = \begin{bmatrix} 1 & 0 & 2 & 0 & 3 & 0 \\ 0 & 1 & 0 & 2 & 0 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The optimal weight vector  $\mathbf{w}^*$  is given by:

$$\mathbf{w}^* = \phi(X)[0.1, 2, 3.9, 5, 6, 8]^T$$

where  $\phi$  is transformation mapping corresponding to the given kernel. What will be the prediction for the data point  $[0, 0, 1]^T$ ?

**Response Type :** Numeric

**Evaluation Required For SA :** Yes

**Show Word Count :** Yes

**Answers Type :** Equal

**Text Areas :** PlainText

**Possible Answers :**

100

**Sub-Section Number :** 6

**Sub-Section Id :** 64065384985

**Question Shuffling Allowed :** Yes

**Is Section Default? :** null

**Question Number : 172 Question Id : 640653588688 Question Type : SA Calculator : None**

**Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 4**

Question Label : Short Answer Question

Consider a ridge regression model with the loss  $L(\mathbf{w}) = \|X^T \mathbf{w} - \mathbf{y}\|^2 + \lambda \|\mathbf{w}\|^2$  is trained on a given dataset with  $\lambda = 0.1, 0, 1, 10, 100$ . Which of the following value of  $\lambda$  is more likely to underfit the model?

**Response Type :** Numeric

**Evaluation Required For SA :** Yes

**Show Word Count :** Yes

**Answers Type :** Equal

**Text Areas :** PlainText

**Possible Answers :**

100

**Question Number : 173 Question Id : 640653588689 Question Type : SA Calculator : None**

**Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 4**

Question Label : Short Answer Question



Consider the following data set:

$$X = [8, 6, 10]$$

Assuming a ridge penalty  $\lambda = 100$ , what will be the value of  $\frac{\hat{w}_{ridge}}{\hat{w}_{MLE}}$ ?

Here  $\hat{w}_{ridge}$  and  $\hat{w}_{MLE}$  are the Ridge and MLE estimates of the weight vectors, respectively.

Assume that the label vector  $y$  of shape  $(3, 1)$  is known. Enter your answer correct to two decimal places.

**Response Type :** Numeric

**Evaluation Required For SA :** Yes

**Show Word Count :** Yes

**Answers Type :** Range

**Text Areas :** PlainText

**Possible Answers :**

0.65 to 0.70

**Sub-Section Number :** 7

**Sub-Section Id :** 64065384986

**Question Shuffling Allowed :** Yes

**Is Section Default? :** null

**Question Number :** 174 **Question Id :** 640653588690 **Question Type :** SA **Calculator :** None

**Response Time :** N.A **Think Time :** N.A **Minimum Instruction Time :** 0

**Correct Marks :** 2.5

**Question Label :** Short Answer Question

A binary classification dataset contains only one feature and the data points given the label follow the Gaussian distributions whose means and variances are already estimated as:

$$x|(y = 0) \sim N(0, 1)$$

$$x|(y = 1) \sim N(2, 2)$$

What will be the prediction for the point  $x = 1$ ? Assume that  $\hat{p}$ , an estimate for  $P(y = 1)$ , is 0.5.

**Response Type :** Numeric

**Evaluation Required For SA :** Yes

**Show Word Count :** Yes



**Answers Type :** Equal

**Text Areas :** PlainText

**Possible Answers :**

0

**Sub-Section Number :** 8

**Sub-Section Id :** 64065384987

**Question Shuffling Allowed :** No

**Is Section Default? :** null

**Question Id : 640653588691 Question Type : COMPREHENSION Sub Question Shuffling Allowed : No Group Comprehension Questions : No Question Pattern Type : NonMatrix Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Question Numbers : (175 to 176)**

Question Label : Comprehension

Consider a binary classification problem and a decision tree that is being trained to classify the points. In one of the internal nodes in this tree, 75% of the data-points belong to one of the two classes and the rest belong to the other class. You are not given the information about which class is more numerous in this node.

Based on the above data, answer the given subquestions.

**Sub questions**

**Question Number : 175 Question Id : 640653588692 Question Type : MCQ Is Question Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 1.5**

Question Label : Multiple Choice Question

Do you have enough information to find the entropy of this node?

**Options :**

6406531963463. ✓ Yes

6406531963464.

✖ No

**Question Number : 176 Question Id : 640653588693 Question Type : SA Calculator : None**

**Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 3**

Question Label : Short Answer Question

If the answer to the previous questions is "Yes", find the entropy of the node. Use  $\log_2$  and enter your answer correct to three decimal places.

If the answer to the previous question is "No", enter -1 as your answer.

**Response Type : Numeric**

**Evaluation Required For SA : Yes**

**Show Word Count : Yes**

**Answers Type : Range**

**Text Areas : PlainText**

**Possible Answers :**

0.79 to 0.83

**Sub-Section Number :** 9

**Sub-Section Id :** 64065384988

**Question Shuffling Allowed :** No

**Is Section Default? :** null

**Question Id : 640653588694 Question Type : COMPREHENSION Sub Question Shuffling**

**Allowed : No Group Comprehension Questions : No Question Pattern Type : NonMatrix**

**Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Question Numbers : (177 to 179)**

Question Label : Comprehension

Consider a probability distribution over  $(X, y)$  where features are one-dimensional and  $y \in \{+1, -1\}$ . Let  $X|(y = 1)$  follow a uniform distribution over  $[0, 4]$  and  $X|(y = -1)$  follows a uniform distribution over  $[2, 4]$ .

Based on the above data, answer the given subquestions.

### Sub questions

**Question Number : 177 Question Id : 640653588695 Question Type : SA Calculator : None**

**Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 2**

Question Label : Short Answer Question

If  $p = P(y = 1)$  is estimated to be 0.4, what will be the prediction for the point  $x = 3$  using the Bayes classifier? Enter 1 or -1.

**Response Type : Numeric**

**Evaluation Required For SA : Yes**

**Show Word Count : Yes**

**Answers Type : Equal**

**Text Areas : PlainText**

**Possible Answers :**

-1

**Question Number : 178 Question Id : 640653588696 Question Type : MCQ Is Question**

**Mandatory : No Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 2**

Question Label : Multiple Choice Question

Let  $x = 2$  and let  $\hat{p}$  be the estimate for  $p = P(y = 1)$ . Find conditions on  $\hat{p}$  such that the Bayes classifier predicts 1 for this  $x$ . Consider that the tie-breaker is predicted in class 1.

**Options :**

6406531963467. ✖  $\hat{p} \leq \frac{1}{4}$

6406531963468. ✖  $\hat{p} \geq \frac{1}{4}$

6406531963469. ✖  $\hat{p} \leq \frac{2}{3}$

6406531963470. ✔  $\hat{p} \geq \frac{2}{3}$

**Question Number : 179 Question Id : 640653588697 Question Type : SA Calculator : None**

**Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Correct Marks : 2**

Question Label : Short Answer Question

If  $p = P(y = 1)$  is estimated to be 0.5 using MLE on a given training dataset, what will be the training error of the Bayes classifier for this problem?

**Response Type : Numeric**

**Evaluation Required For SA : Yes**

**Show Word Count : Yes**

**Answers Type : Equal**

**Text Areas : PlainText**

**Possible Answers :**

0.5

**Sub-Section Number :** 10

**Sub-Section Id :** 64065384989

**Question Shuffling Allowed :** No

**Is Section Default? :** null

**Question Id : 640653588698 Question Type : COMPREHENSION Sub Question Shuffling Allowed : No Group Comprehension Questions : No Question Pattern Type : NonMatrix Calculator : None Response Time : N.A Think Time : N.A Minimum Instruction Time : 0**

**Question Numbers : (180 to 181)**

Question Label : Comprehension

Consider a naive Bayes model is trained on the following data matrix  $X$  of shape  $(d, n)$  and corresponding label vector  $y$ :

$$X = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \end{bmatrix} \quad y = [1 \ 0 \ 1 \ 0]^T$$

Assume that  $\hat{p}$  and  $\hat{p}_j^{y_i}$  are estimates for  $P(y = 1)$  and  $P(f_j = 1|y = y_i)$ , respectively. Here,  $f_i$ ;  $i = 1, 2, 3$  is the  $i^{\text{th}}$  feature. These parameters are estimated using MLE. Do not apply any smoothing on the dataset.

Based on the above data, answer the given subquestions.

**Sub questions****Question Number : 180 Question Id : 640653588699 Question Type : SA Calculator : None****Response Time : N.A Think Time : N.A Minimum Instruction Time : 0****Correct Marks : 2**

Question Label : Short Answer Question

Calculate the value of  $\hat{p}_2^0$ .**Response Type : Numeric****Evaluation Required For SA : Yes****Show Word Count : Yes****Answers Type : Equal****Text Areas : PlainText****Possible Answers :**

0.5

**Question Number : 181 Question Id : 640653588700 Question Type : SA Calculator : None****Response Time : N.A Think Time : N.A Minimum Instruction Time : 0****Correct Marks : 2**

Question Label : Short Answer Question

Calculate the value of  $\hat{p}_2^1$ .

**Response Type :** Numeric

**Evaluation Required For SA :** Yes

**Show Word Count :** Yes

**Answers Type :** Equal

**Text Areas :** PlainText

**Possible Answers :**

0

## Java

<b>Section Id :</b>	64065339804
<b>Section Number :</b>	11
<b>Section type :</b>	Online
<b>Mandatory or Optional :</b>	Mandatory
<b>Number of Questions :</b>	16
<b>Number of Questions to be attempted :</b>	16
<b>Section Marks :</b>	50
<b>Display Number Panel :</b>	Yes
<b>Group All Questions :</b>	No
<b>Enable Mark as Answered Mark for Review and Clear Response :</b>	Yes
<b>Maximum Instruction Time :</b>	0
<b>Sub-Section Number :</b>	1
<b>Sub-Section Id :</b>	64065384990
<b>Question Shuffling Allowed :</b>	No
<b>Is Section Default? :</b>	null

**Question Number :** 182 **Question Id :** 640653588701 **Question Type :** MCQ **Is Question**

**Mandatory :** No **Calculator :** None **Response Time :** N.A **Think Time :** N.A **Minimum Instruction**